

An inequality for the h -invariant in instanton Floer theory

Kim A. Frøyshov*

4 March 2003

Abstract

Given a smooth, compact, oriented 4-manifold X with a homology sphere Y as boundary and $b_2^+(X) = 1$, and given an embedded surface $\Sigma \subset X$ of self-intersection 1, we prove an inequality relating $h(Y)$, the genus of Σ , and a certain invariant of the orthogonal complement of $[\Sigma]$ in the intersection form of X .

1 Introduction

In [10] we defined a surjective group homomorphism from the homology cobordism group of oriented (integral) homology 3-spheres to the integers,

$$h : \theta_3^H \rightarrow \mathbb{Z},$$

using instanton Floer theory. The main purpose of this paper is to establish the following property of this invariant.

Theorem 1 *Let X be a smooth, compact, oriented 4-manifold with a homology sphere Y as boundary and with $b_2^+(X) = 1$. Let $\Sigma \subset X$ be a closed surface of genus g and self-intersection $\Sigma \cdot \Sigma = 1$, and let $\mathcal{L} \subset H^2(X; \mathbb{Z})/\text{torsion}$ be the sublattice consisting of all vectors which vanish on $[\Sigma]$. Then the intersection form of X restricts to a unimodular negative definite form on $\mathcal{L}$, and we have*

$$h(Y) + \lceil g/2 \rceil \geq e(\mathcal{L}),$$

where $e(\mathcal{L})$ is a non-negative integer which depends only on $\mathcal{L}$. Moreover,

*Partially supported by a post-doctoral grant from the Norwegian Research Council, and by NSF grant DMS-9971731.

- $e(\mathcal{L}) = 0$ precisely when $\mathcal{L}$ is diagonal
- $e(-kE_8 \oplus \mathcal{L}) = k$ if $\mathcal{L}$ is diagonal and $k \geq 0$.

By “surface in X ” we mean a 2-dimensional compact, oriented, connected smooth submanifold of X . $\lceil x \rceil$ denotes the smallest integer $\geq x$.

See Section 2 for a slightly more general statement and the definition of $e(\mathcal{L})$. To compute the term $\lceil g/2 \rceil$ in the inequality we rely on Muñoz’ description [14] of the ring structure of the Floer cohomology of the $\mathrm{SO}(3)$ bundle $E \rightarrow S^1 \times \Sigma$ where E is the pull-back of the non-trivial $\mathrm{SO}(3)$ bundle over Σ .

Note that if Z is a negative definite 4-manifold with a homology sphere Y as boundary then one can apply the theorem to $X = Z \# \mathbb{CP}^2$ with $g = 0$. When $Y = S^3$ one recovers Donaldson’s diagonalization theorem [3, 4].

As another example, we obtain bounds on how much h may change under ± 1 surgery on knots:

Corollary 1 *Let Y be an oriented homology 3-sphere and γ a knot in Y of slice genus g . If $Y_{\gamma, -1}$ is the result of -1 surgery on γ then*

$$0 \leq h(Y_{\gamma, -1}) - h(Y) \leq \lceil g/2 \rceil.$$

Here the slice genus may be defined as the smallest non-negative integer g for which there exists a smooth rational homology cobordism W from Y to some rational homology sphere Y' and a genus g surface $\Sigma \subset W$ such that $\partial\Sigma = \gamma$. (We do not know whether this definition agrees with the usual one for $Y = S^3$.)

To deduce the corollary from the theorem, let Z be the surgery cobordism from Y to $Y_{\gamma, -1}$ and set $W' = W \cup_{Y'} \overline{W} \cup_Y Z$, which is a cobordism from Y to $Y_{\gamma, -1}$. By attaching a suitable 1-handle to W' we obtain a smooth, compact, oriented 4-manifold X with boundary $\overline{Y} \# Y_{\gamma, -1}$ and such that X contains a closed surface of genus g and self-intersection -1 representing a generator of $H_2(X; \mathbb{Z})/\text{torsion}$. Now apply Theorem 1 to X with both orientations and the Corollary follows, since h is additive under connected sums.

If γ is the (p, q) torus knot in S^3 , where p, q are mutually prime integers ≥ 2 , then $Y_{\gamma, -1}$ is diffeomorphic to the Brieskorn sphere $\Sigma(p, q, pq - 1)$. In this case one can also apply Theorem 1 to the minimal resolution of the corresponding Brieskorn singularity to get a lower bound on $h(Y_{\gamma, -1})$. In general, this lower bound does not coincide with the upper bound given by

Corollary 1. However, we will show in the next section that they do coincide when $p = 2$, allowing us to compute

$$h(\Sigma(2, 2k - 1, 4k - 3)) = \lfloor k/2 \rfloor$$

for $k \geq 2$, where $\lfloor x \rfloor$ is the largest integer $\leq x$.

Contents

1	Introduction	1
2	The general inequality	4
3	Outline of proof	6
4	Parametrized moduli spaces	8
4.1	Orientations	8
4.2	Local structure near reducibles	10
5	The family of metrics	13
6	Reducibles	14
7	Cutting down parametrized moduli spaces	17
8	Main mechanism of proof	19
9	Boundary points of $\widehat{\mathcal{M}}$	21
10	Ends of $\widehat{\mathcal{M}}$	22
11	μ-classes of loops	25
A	Determinant line bundles	28
A.1	Determinant lines	28
A.2	Determinant line bundles	30
A.3	Isomorphisms between determinant line bundles	32
B	On projectively flat $U(2)$ connections over 3-manifolds	33

2 The general inequality

We will now state the main result of this paper, which is more general than Theorem 1. We first recall some definitions from [10].

By a *lattice* we shall mean a finitely generated free abelian group $\mathcal{L}$ with a non-degenerate symmetric bilinear form $\mathcal{L} \times \mathcal{L} \rightarrow \mathbb{Z}$. The dual lattice $\text{Hom}(\mathcal{L}, \mathbb{Z})$ will be denoted $\mathcal{L}^\#$.

Definition 1 *Let $\mathcal{L}$ be a (positive or negative) definite lattice. A vector $w \in \mathcal{L}$ is called extremal if $|w^2| \leq |z^2|$ for all $z \in w + 2\mathcal{L}$. If $w \in \mathcal{L}$, $a \in \mathcal{L}^\#$, and m is a non-negative integer satisfying $w^2 \equiv m \pmod{2}$ set*

$$\eta(\mathcal{L}, w, a, m) = \sum_z (-1)^{((z+w)/2)^2} (a \cdot z)^m,$$

where the sum is taken over all $z \in w + 2\mathcal{L}$ such that $z^2 = w^2$. If $m = 0$ then we interpret $(a \cdot z)^m = 1$ and write $\eta(\mathcal{L}, w) = \eta(\mathcal{L}, w, a, m)$.

Notice that $(-1)^{((z+w)/2)^2} (a \cdot z)^m$ remains the same when z is replaced with $-z$. Therefore, if $w \neq 0$ then this definition of η differs from that in [10] by a factor of 2.

Theorem 2 *Let X be a smooth, compact, oriented 4-manifold with a homology sphere Y as boundary and with $b_2^+(X) \geq 1$. For $i = 1, \dots, b_2^+(X)$ let $\Sigma_i \subset X$ be a closed surface of self-intersection 1, such that $\Sigma_i \cap \Sigma_j = \emptyset$ for $i \neq j$. Set $g = \text{genus}(\Sigma_1)$ and suppose $\text{genus}(\Sigma_i) = 1$ for $i \geq 2$. Let $\mathcal{L} \subset H^2(X; \mathbb{Z})/\text{torsion}$ be the sublattice consisting of all vectors which vanish on all the classes $[\Sigma_i]$. Thus, $\mathcal{L}$ is a unimodular, negative definite form. Let $w \in \mathcal{L}$ be an extremal vector, let $a \in H_2(X; \mathbb{Z})$, and let m be a non-negative integer such that $w^2 \equiv m \pmod{2}$ and $\eta(\mathcal{L}, w, a, m) \neq 0$. Then*

$$h(Y) + \lceil g/2 \rceil + b_2^+(X) - 1 \geq (|w^2| - m)/4.$$

The author does not see any natural generalization of the theorem in which the roles of the Σ_i are symmetric. (See Section 10 for an explanation of this.)

Definition 2 *For any definite lattice $\mathcal{L}$ let $e(\mathcal{L})$ be the supremum of the set of all integers $\lceil (|w^2| - m)/4 \rceil$ where w is any extremal vector in $\mathcal{L}$ and m any non-negative integer such that (i) $w^2 \equiv m \pmod{2}$, and (ii) $\eta(\mathcal{L}, w, a, m) \neq 0$ for some $a \in \mathcal{L}^\#$.*

With this definition of $e(\mathcal{L})$, the inequality in Theorem 1 follows from Theorem 2. Since every unimodular lattice can be realized as the intersection form of a smooth, compact, oriented 4-manifold with a homology sphere as boundary, Theorem 2 implies that $e(\mathcal{L})$ is finite for any unimodular definite lattice $\mathcal{L}$.

Applying Theorem 2 to $(\mathbb{CP}^2 \# k \overline{\mathbb{CP}^2}) \setminus (\text{open ball})$ we see that $e(\mathcal{L}) = 0$ when $\mathcal{L}$ is diagonal. (Of course, one can also verify this directly.) If $\mathcal{L}$ is not diagonal then $e(\mathcal{L}) > 0$ by the proof of [10, Corollary 2]. In a similar fashion one can show that $e(-kE_8 \oplus \mathcal{L}) = k$ if $\mathcal{L}$ is diagonal (cfr. the proof of Proposition 2 below).

Before embarking on the proof of Theorem 2 we apply it to compute h in some examples.

Proposition 1 $h(\Sigma(2, 2k - 1, 4k - 3)) = \lfloor k/2 \rfloor$ for $k \geq 2$.

Proof. Set $Y_k = \Sigma(2, 2k - 1, 4k - 3)$. The minimal resolution of the corresponding Brieskorn singularity has intersection form isomorphic to

$$\Gamma_{4k} = \{ \sum x_i e_i \in \mathbb{R}^{4k} \mid \sum x_i \in 2\mathbb{Z}; 2x_i \in \mathbb{Z}; x_i - x_j \in \mathbb{Z} \}, \quad (1)$$

where $\{e_i\}$ is an orthonormal basis for $\mathbb{R}^{4k}$. This is an even form precisely when k is even. Set $\ell = \lfloor k/2 \rfloor$. It is easy to see that $w = \sum_{i=1}^{4\ell} e_i$ is an extremal vector in Γ_{4k} with $\eta(\Gamma_{4k}, w) \neq 0$. Hence $h(Y_k) \geq |w|^2/4 = \ell$ by Theorem 2. On the other hand, Y_k is also -1 surgery on the $(2, 2k - 1)$ torus knot, which has genus $k - 1$. Since $\lceil (k - 1)/2 \rceil = \lfloor k/2 \rfloor = \ell$, Corollary 1 gives $h(Y_k) \leq \ell$. $\square$

The first step in our proof of Theorem 2 is the following proposition, which uses the additivity of h to reduce the theorem to the case $h(Y) = 0$, at the expense of perhaps increasing $b_2^+(X)$. In the simplest case $b_2^+(X) = 1$ it is natural to ask for an alternative proof which only uses 4-manifolds with $b_2^+ = 1$. Unfortunately, such a proof would seem to require an extensive discussion of bubbles, which we will not take up here.

Proposition 2 *If Theorem 2 holds when $h(Y) = 0$ and $b_1(X) = 0$ then it holds in general.*

Proof. Performing surgery on a set of loops in X representing a basis for the free part of $H_1(X; \mathbb{Z})$ yields a 4-manifold with $b_1 = 0$ but with the same intersection form and the same torsion in $H_1(\cdot; \mathbb{Z})$. Since the loops can be chosen disjoint from the surfaces Σ_i , it suffices to prove the Theorem when $b_1(X) = 0$.

Let X_1 denote the (negative definite) E_8 -manifold with boundary $S = \Sigma(2, 3, 5)$. Let X_2 denote the oriented 4-manifold described by the right-handed trefoil in S^3 with framing $+1$. Then $\partial X_2 = \overline{S}$ and X_2 contains an embedded torus of self-intersection $+1$ which represents a generator of $H_2(X_2; \mathbb{Z}) = \mathbb{Z}$.

For any pair Z_1, Z_2 of oriented 4-manifolds with non-empty and connected boundaries let $Z_1 \#_{\partial} Z_2$ denote their “boundary sum”, formed by joining the boundaries of Z_1 and Z_2 by a 1-handle (respecting the orientations of Z_1 and Z_2).

If $h(Y) = -k < 0$ we replace X in the Theorem by the k -fold boundary connected sum $X' = X \#_{\partial} kX_1$. Then $\partial X' = Y \# kS$ has $h = 0$. Also, replace w by $w' = w + \sum_{j=1}^k q_j$, where q_j is supported on the j 'th copy of X_1 and is given by $q_j = e_1 + e_2 + e_3 + e_4 \in E_8 = \Gamma_8$, in the notation of (1). Since $\eta(E_8, e_1 + e_2 + e_3 + e_4) = 16$, we have altered $\eta(\mathcal{L}, w, a, m)$ by a factor of 16^k . This reduces the Theorem to the case $h(Y) \geq 0$.

If $h(Y) = k > 0$ replace X by $X' = X \#_{\partial} kX_2$, so $\partial X' = Y \# k\overline{S}$, which has $h = 0$. The k copies of X_2 provide k embedded tori in X' with self-intersection 1. $\square$

3 Outline of proof

It remains to prove Theorem 2 in the case $h(Y) = 0 = b_1(X)$. In this outline we will assume $Y = S^3$, since the proof in the case $h(Y) = 0$ is almost the same. We can then just as well assume X is closed. We will also take $b_2^+(X) = 1$, $H_1(X; \mathbb{Z}) = 0$, and $m = 0$. What we need to prove then is that if $w \in \mathcal{L}$ is an extremal vector such that w^2 is even and $\eta(\mathcal{L}, w) \neq 0$ then

$$2q \geq n + 1,$$

where $q = \lceil g/2 \rceil$ and $n + 1 = |w^2|/2$.

Let N be a closed tubular neighbourhood of Σ and W the result of replacing $N \subset X$ by $V = N \# \overline{\mathbb{CP}^2}$. After choosing orientations of Σ and $S = \mathbb{CP}^1 \subset \mathbb{CP}^2$ we form two internal connected sums $\Sigma^{\pm} \subset V$ of Σ and S , in one case preserving orientations and in the other case reversing them. Thus $\Sigma^{\pm}$ has self-intersection number 0. We construct a smooth 1-parameter family of metrics $g(t)$ on W , independent of t outside V , as follows: First choose an initial metric in which a small tubular neighbourhood of $C = \partial V$ is isometric to $[0, T] \times C$, where $T \gg 0$. Then stretch this initial metric along the link of Σ^+ (for $t > 1$) and along the link of Σ^- (for $t < -1$). More

precisely, when $t > 1$ then W should contain a cylinder $[0, t] \times S^1 \times \Sigma^+$, and similarly for $t < -1$.

Let $s \in H^2(W; \mathbb{Z})$ be the Poincaré dual of $[S]$. Choose an integral lift w' of w and for any integer k let $E_k \rightarrow W$ be the $U(2)$ bundle with $c_1 = w' + s$ and $c_2 = k$. For each t let $M_{k,t}$ be the moduli space of projectively $g(t)$ anti-self-dual connections in E_k . Let $\mathcal{M}_k$ denote the disjoint union of the $M_{k,t}$, $t \in \mathbb{R}$. For suitable, small perturbations of the ASD equations, the irreducible part $\mathcal{M}_k^*$ of $\mathcal{M}_k$ will be a smooth manifold of dimension

$$\dim \mathcal{M}_k^* = 8k + 4n + 1.$$

In Section 6 we will show that $\mathcal{M}_k$ contains no reducibles for $k < 0$, while $\mathcal{M}_0$ contains a finite number of reducibles (for a generic choice of initial metric). Moreover, one can describe explicitly, in terms of $\mathcal{L}$ and w only, the splittings of E_0 (into complex line bundles) that correspond to reducibles in $\mathcal{M}_0$. To each such splitting there is associated a “degree” (always ± 1) which measures the number of times (counted with sign) that this splitting occurs. In addition to the degree, each splitting also comes with a sign which measures whether the overall orientation of a certain determinant line bundle over the orbit space of connections in E_0 agrees with the “complex orientation” at the corresponding reducible points. The sum of (degree)·(sign) over all the splittings is equal to $2\eta(\mathcal{L}, w)$.

Let $\mathcal{M}'_0 \subset \mathcal{M}_0$ be the result of removing from $\mathcal{M}_0$ a small open neighbourhood about each reducible point. For $k < 0$ set $\mathcal{M}'_k = \mathcal{M}_k$. Recall that to any base-point in W one can associate a principal $SO(3)$ bundle $\mathbb{E} \rightarrow \mathcal{M}'_k$ called the base-point fibration. For $j \geq 0$ and any subset S of $\mathcal{M}'_k$ let $x^j \cdot S$ denote the intersection of S with j generic geometric representatives for $-\frac{1}{4}p_1(\mathbb{E})$, in the sense of [12]. These representatives should depend only on the restriction of elements of S to suitable compact subsets of $W \setminus V$ where the metric does not vary with t .

We also need a variant of this construction where the location of the base-point depends on t ; in this case the result is denoted $x^j_1 \cdot S$. To make this more precise: for $\pm t \gg 0$ the geometric representatives should be defined through restriction to subsets of some tubular neighbourhood of $\Sigma^\pm$ where the metric does not vary, while for intermediate values of t we interpolate, in a certain sense.

Now suppose the theorem does not hold, ie that $2q \leq n$, and set

$$\widehat{\mathcal{M}} = \text{1-dimensional part of } x^{n-2q} \cdot (x_1^2 - 4)^q \cdot \sum_{k \leq 0} \mathcal{M}'_k.$$

This is a formal linear combination of oriented 1-manifolds with boundary. The number of boundary points of $\widehat{\mathcal{M}}$, counted with multiplicity, equals $\pm 4^{-n} \eta(\mathcal{L}, w)$. On the other hand, it follows from Muñoz' description of the ring structure of the Floer cohomology of $S^1 \times \Sigma^\pm$ that the number of ends of $\widehat{\mathcal{M}}$, counted with multiplicity, is zero. This contradiction proves the theorem in the case considered.

4 Parametrized moduli spaces

In this section we study instanton moduli spaces over a 4-manifold W with tubular ends, parametrized by a family of tubular end metrics. We are interested in questions of orientations and reducibles.

4.1 Orientations

Let W be an oriented Riemannian 4-manifold with tubular ends $\mathbb{R}_+ \times Y_j$, $j = 1, \dots, r$, so $W \setminus \cup_j (\mathbb{R}_+ \times Y_j)$ is compact. Let $E \rightarrow W$ be a $U(2)$ bundle. We denote by $\mathfrak{g}_E$ the bundle of Lie algebras associated to E and the adjoint representation of $U(2)$ on its Lie algebra $\mathfrak{u}(2)$, and by $\mathfrak{g}'_E$ the sub-bundle corresponding to the sub-algebra $\mathfrak{su}(2) \subset \mathfrak{u}(2)$.

Choose an isomorphism $E|_{\mathbb{R}_+ \times Y_j} = \mathbb{R} \times E_j$, where E_j is a $U(2)$ bundle over Y_j . Let α_j be a connection in E_j such that the induced connection in $\mathfrak{g}'_{E_j}$ is non-degenerate flat. (As usual, 'flat connections' will in practice mean critical points of the perturbed Chern-Simons functional as in [8].) Let $\sigma_j \geq 0$ be small, and $\sigma_j > 0$ if α_j is reducible. Fix an even integer $p > 4$. Choose a smooth connection A_0 in E which agrees with the pull-back of α_j over the j 'th end, and define the Sobolev space of connections

$$\mathcal{A} = \mathcal{A}(E, \alpha) = \{A_0 + a \mid a \in L_1^{p,\sigma}(T^*W \otimes \mathfrak{g}'_E)\}.$$

Here $L_k^{p,\sigma}$ is the space of sections s such that $(\nabla_{A_0})^j(e^w s) \in L^p$ weakly for $0 \leq j \leq k$, where $w : W \rightarrow \mathbb{R}$ is any smooth function with $w(t, y) = e^{\sigma_j t}$ for $(t, y) \in \mathbb{R}_+ \times Y_j$. As explained in [6] there is a Banach Lie group $\mathcal{G}$ consisting of $L_{2,\text{loc}}^p$ gauge transformations, such that $\mathcal{G}$ acts smoothly on $\mathcal{A}$ and the following holds: If $A, B \in \mathcal{A}$ and there is an $L_{2,\text{loc}}^p$ gauge transformation u such that $u(A) = B$ then $u \in \mathcal{G}$.

There is a real determinant line bundle $\det(\delta)$ over $\mathcal{B} = \mathcal{A}/\mathcal{G}$ associated to the family of Fredholm operators

$$\delta_A = d_A^* + d_A^+ : L_1^{p,\sigma} \rightarrow L^{p,\sigma},$$

see [7, 6] and Appendix A. It is proved in [6] that $\det(\delta)$ is orientable. An orientation of $\det(\delta)$ defines an orientation of the regular part of the instanton moduli space $M^* \subset \mathcal{B}^*$ cut out by the equation $F_0^+(A) = 0$, where $F_0(A)$ is the curvature of the connection that A induces in $\mathfrak{g}'_E$. As usual, $\mathcal{B}^*$ denotes the irreducible part of $\mathcal{B}$, etc. As for the choice of orientation, the important thing for us will be that orientations of the various moduli spaces involved be chosen compatible with gluing maps, and this can be done at least in the situations we will consider.

More generally, let $g(t)$ be a smooth family of Riemannian metrics on W , where t runs through some parameter space $\mathbb{R}^b$, such that $g(t)$ is constant in t outside some compact set in W . In the following let Λ^j be the bundle of j -forms and Λ_t^+ the bundle of $g(t)$ self-dual 2-forms over W . There is then a parametrized moduli space

$$\mathcal{M} \subset \mathcal{B} \times \mathbb{R}^b$$

consisting of all pairs $([A], t)$ satisfying

$$P_t^+ F_0(A) = 0, \quad (2)$$

where $P_t : \Lambda^2 \rightarrow \Lambda_t^+$. Generically, the irreducible part of this moduli space, $\mathcal{M}^*$, will be a smooth manifold. (In the situations encountered in this paper it will suffice to start with the usual translationary invariant perturbations over the ends of W , given by perturbations of the Chern-Simons functional as in [8], and add further perturbations defined in terms of holonomy along a finite number of thickened loops in W . In general, perturbations should be small so that one can control reducibles.)

To orient $\mathcal{M}^*$ we note that if t is sufficiently close to a reference point τ then (2) is equivalent to

$$P_\tau^+ P_t^+ F_0(A) = 0.$$

The derivative of $(A, t) \mapsto P_\tau^+ P_t^+ F_0(A)$ at (A, τ) is the operator

$$\begin{aligned} T_{A,\tau} : L_1^{p,\sigma}(\Lambda^1 \otimes \mathfrak{g}'_E) \times \mathbb{R}^b &\rightarrow L^{p,\sigma}(\Lambda_\tau^+ \otimes \mathfrak{g}'_E) \\ (a, x) &\mapsto P_\tau^+ d_A a + S_{A,\tau} x, \end{aligned}$$

where for any $(A, \tau) \in \mathcal{A} \times \mathbb{R}^b$ we define

$$S_{A,\tau} x = \left. \frac{\partial}{\partial s} \right|_0 P_\tau^+ P_{\tau+sx}^+ F_0(A).$$

Differentiating the equation $P_\tau^+ P_\tau^- = 0$ we obtain

$$\frac{\partial}{\partial s} (P_{\tau+sx}^+ P_\tau^- - P_\tau^+ P_{\tau+sx}^-) = 0,$$

hence

$$S_{A,\tau}x = \frac{\partial}{\partial s} \Big|_0 P_{\tau+sx}^+ F_0(A) \quad \text{if } ([A], \tau) \in \mathcal{M}.$$

A point $([A], t) \in \mathcal{M}$ is called *regular* if $T_{A,t}$ is surjective. The tangent space of $\mathcal{M}$ at an irreducible regular point $([A], t)$ can be identified with the kernel of the operator

$$\tilde{\delta}_{A,t} = \delta_{A,t} \oplus S_{A,t} : L_1^{p,\sigma}(\Lambda^1 \otimes \mathfrak{g}'_E) \oplus \mathbb{R}^b \rightarrow L^{p,\sigma}(\mathfrak{g}'_E) \oplus L^{p,\sigma}(\Lambda_t^+ \otimes \mathfrak{g}'_E),$$

where $\delta_{A,t}$ denotes the operator δ_A in the metric $g(t)$. What we are seeking, therefore, is an orientation of the determinant line bundle $\det(\tilde{\delta})$ over $\mathcal{B} \times \mathbb{R}^b$ associated to the family of Fredholm operators $\tilde{\delta}_{A,t}$. But as explained in Subsection A.3 there is a canonical isomorphism

$$\gamma : \det(\delta) \rightarrow \det(\tilde{\delta}) \otimes (\det(\mathbb{R}^b))^*. \quad (3)$$

Given an orientation of $\det(\delta)$, this orients $\det(\tilde{\delta})$, hence $\mathcal{M}^*$.

4.2 Local structure near reducibles

We now assume, for simplicity, that each Y_j is a rational homology sphere, and that $b_1(X) = 0$. More importantly, we take $b = b^+$, where $b^+ = b_2^+(W)$.

Let $D = ([A], t) \in \mathcal{M}$ be a reducible point, so A respects some splitting of E into complex line bundles, $E = L_1 \oplus L_2$. Then $\mathfrak{g}'_E = \underline{\mathbb{R}} \oplus K$, where $K = L_1 \otimes \bar{L}_2$ and $\underline{\mathbb{R}} = W \times \mathbb{R}$. Here the constant section $\mathbf{1}$ of $\underline{\mathbb{R}}$ acts as $2i$ on K . Note that changing the order of L_1, L_2 changes the orientation of K , hence many signs in what follows.

Let B denote the connection that A induces in K . Noting that $F_0(A)$ takes values in $\underline{\mathbb{R}}$, we can identify $F_0(A) = -2iF(B)$, which is an anti-self-dual, closed L^2 2-form representing $4\pi c_1(K)$.

Let $\mathcal{H}_t^+$ be the space of self-dual, closed, L^2 2-forms on $(W, g(t))$. By [1, Proposition 4.9] we can identify $\mathcal{H}_t^+$ with a subspace of $H^2(W)$.

The operator

$$P_t^+ d_A : L_1^{p,\sigma}(\Lambda^1 \otimes \mathfrak{g}'_E) \rightarrow L^{p,\sigma}(\Lambda_t^+ \otimes \mathfrak{g}'_E)$$

is the sum of two operators: one with values in $\underline{\mathbb{R}}$, which is the usual $P_t^+ d$ operator, and another with values in K , which we call $P_t^+ d_B$. Note that the dual of the cokernel of $P_t^+ d$ is $\mathcal{H}_t^+$, and the map $\mathbb{R}^{b^+} \rightarrow \text{coker}(\delta_{A,t})$ defined by $S_{A,t}$ takes values in $(\mathcal{H}_t^+)^*$. Let $R_D : \mathbb{R}^{b^+} \rightarrow (\mathcal{H}_t^+)^*$ denote this operator.

Observation 1 *D is a regular point of $\mathcal{M}$ if and only if the following two conditions hold:*

- (i) $P_t^+ d_B : L_1^{p,\sigma}(\Lambda^1 \otimes K) \rightarrow L^{p,\sigma}(\Lambda_t^+ \otimes K)$ is surjective.
- (ii) $R_D : \mathbb{R}^{b^+} \rightarrow (\mathcal{H}_t^+)^*$ is an isomorphism.

Note that if the bundle K is non-trivial then, in the metric $g(t)$, d_B^+ is surjective if and only if $d_B^* + d_B^+$ is surjective, in which case the latter operator has non-negative index. So if this index is negative then D cannot be a regular point of $\mathcal{M}$. On the other hand, if the index is non-negative (which will be the case in our applications) then as explained in [3] there is a simple local perturbation of the anti-self-duality equation near $D \in \mathcal{B} \times \mathbb{R}^{b^+}$ such that D solves the perturbed equation and such that the perturbed analogue of $P_t^+ d_B$ is surjective. (Ideally, one should look for a generalization of Freed and Uhlenbeck's theorem to our situation, but we will not pursue this here.)

We will now give a more concrete description of the map R_D . Let $\Sigma_1, \dots, \Sigma_{b^+} \subset W$ be closed surfaces such that $\Sigma_i \cdot \Sigma_j = 0$ if $i \neq j$ and $\Sigma_i \cdot \Sigma_i > 0$ for all i . Then there is a linear isomorphism

$$\mathcal{H}_t^+ \rightarrow \mathbb{R}^{b^+}, \quad \omega \mapsto \left(\int_{\Sigma_i} \omega \right)_{1 \leq i \leq b^+}.$$

We can therefore define a basis $\{\omega_{i,t}\} \subset \mathcal{H}_t^+$ by the conditions

$$\int_W \omega_{i,t} \wedge \omega_{i,t} = 1; \quad \int_{\Sigma_i} \omega_{i,t} > 0; \quad \int_{\Sigma_i} \omega_{j,t} = 0 \quad \text{for } i \neq j.$$

For any $v \in H^2(W)$ consider the map

$$f_v : \mathbb{R}^{b^+} \rightarrow \mathbb{R}^{b^+}, \quad t \mapsto \left(\int_W \omega_{i,t} \wedge v \right)_{1 \leq i \leq b^+}, \quad (4)$$

where in the integral v is represented by some bounded, closed 2-form. Clearly, $f_v(t) = 0$ if and only if v can be represented by a $g(t)$ anti-self-dual, closed L^2 form.

Proposition 3 *With respect to the basis for $\mathcal{H}_t^+$ defined above we have*

$$R_D = df_v(t),$$

where $v = 4\pi c_1(K)$, and $df_v(t)$ is the derivative of the function f_v at t .

Proof. For every $x \in \mathbb{R}^{b^+}$ we have

$$\begin{aligned}
R_D x \cdot \omega_{i,t} &= \frac{d}{ds} \Big|_0 \int_W P_{t+sx}^+ F_0(A) \wedge \omega_{i,t} \\
&= \frac{d}{ds} \Big|_0 \int_W P_{t+sx}^+ F_0(A) \wedge \omega_{i,t+sx} \\
&= \frac{d}{ds} \Big|_0 \int_W v \wedge \omega_{i,t+sx} \\
&= df_{v,i}(t)x,
\end{aligned}$$

where $f_{v,i}$ is the i 'th component of f_v . $\square$

We conclude this section with a simple result about orientations. Suppose D is a regular point. Then $\ker(\delta_D) = \ker(\tilde{\delta}_D)$ is a complex vector space (recall that we assume $b_1(W) = 0$), and therefore has a canonical orientation. Furthermore,

$$\operatorname{coker}(\tilde{\delta}_D) = \mathbb{R}\mathbf{1}, \quad \operatorname{coker}(\delta_D) = \mathbb{R}\mathbf{1} \oplus (\mathcal{H}_t^+)^*.$$

Given the ordering of the surfaces Σ_i we then obtain orientations of $\det(\delta_D)$ and $\det(\tilde{\delta}_D)$, which we refer to as the “complex orientations”, cfr. [4]. We would like to know how these compare under the isomorphism γ_D in (3). As explained in Subsection A.1 the exact sequence

$$0 \rightarrow \ker(\delta_D) \rightarrow \ker(\tilde{\delta}_D) \rightarrow \mathbb{R}^b \rightarrow \operatorname{coker}(\delta_D) \rightarrow \operatorname{coker}(\tilde{\delta}_D) \rightarrow 0$$

gives rise to a natural isomorphism

$$\gamma'_D : \det(\delta_D) \rightarrow \det(\tilde{\delta}_D) \otimes (\det(\mathbb{R}^b))^*.$$

By Proposition 9 (with $\epsilon_1 = \epsilon_2 = 1$) we have

$$\gamma_D = \gamma'_D. \tag{5}$$

This yields:

Proposition 4 *If D is a regular point then the complex orientations of $\det(\delta_D)$ and $\det(\tilde{\delta}_D)$ agree under the isomorphism γ_D if and only if R_D preserves orientation.*

5 The family of metrics

We can now begin the proof of Theorem 2 in earnest. By Proposition 2 we may assume $b_1(X) = 0$. (The assumption $h(Y) = 0$ will not be used until Section 8.) In this section we construct from X the 4-manifold W that will be the base-manifold for our moduli spaces later. We obtain a $b_2^+(W) + 1$ dimensional family of Riemannian metrics on W by stretching along various hypersurfaces.

To define W , set $b^+ = b_2^+(X)$ and let $\hat{X}$ be the result of adding a half-infinite tube $\mathbb{R}_+ \times Y$ to X . Choose disjoint, compact tubular neighbourhoods N_i of the surfaces Σ_i and let $V_i \approx N_i \# \mathbb{CP}^2$ be the blow-up of N_i at some interior point away from Σ_i . We then define W to be the manifold obtained from $\hat{X}$ by replacing each N_i by V_i . Thus,

$$W \approx \hat{X} \# b^+ \mathbb{CP}^2.$$

Let $S_i \subset \mathbb{CP}^2$ be a sphere representing a generator of $H_2(\mathbb{CP}^2; \mathbb{Z})$. Choose internal connected sums $\Sigma_i^+ = \Sigma_i \# S_i$ and $\Sigma_i^- = \Sigma_i \# \overline{S}_i$ in the interior of V_i .

Let S^1 be the boundary of the closed unit disk $D^2 \subset \mathbb{R}^2$ centered at the origin. For each i choose smooth embeddings

$$q_i^\pm : D^2 \times \Sigma_i^\pm \rightarrow V_i$$

such that $q_i^\pm(0, \cdot)$ is the identity on $\Sigma_i^\pm$. Let $N_i^\pm$ denote the image of the embedding

$$\begin{aligned} \rho_i^\pm : [0, 1] \times S^1 \times \Sigma_i^\pm &\rightarrow V_i \\ (s, x, z) &\mapsto q_i^\pm\left(\frac{1}{2}(s+1)x, z\right). \end{aligned}$$

Set $W' = W \setminus (\cup_i \text{int } V_i)$ and $C = \partial W'$, and let N_C be the image of a smooth embedding

$$\rho_C : [0, 1] \times C \rightarrow W'$$

such that $\rho_C(0, \cdot)$ is the identity on C . Set $W^- = W' \setminus \rho_C([0, 1] \times C)$. Choose Riemannian metrics on C and on $S^1 \times \Sigma_i^\pm$.

Choose a smooth function $\tau : \mathbb{R} \rightarrow \mathbb{R}$ such that $\tau' \geq 0$, $\tau(s) = 0$ for $s \leq \frac{1}{3}$, $\tau(s) = 1$ for $s \geq \frac{2}{3}$. For $r \geq 1$ let $\xi_r : [0, r] \rightarrow [0, 1]$ be the diffeomorphism whose inverse is given by

$$\xi_r^{-1}(s) = s + (r-1)\tau(s).$$

Now choose a smooth family of Riemannian metrics $g(T, t)$ on W , where $T \geq 1$ and $t = (t_1, \dots, t_{b^+}) \in \mathbb{R}^{b^+}$, such that the following holds:

- If N_C and $N_i^\pm$ have the metrics induced by $g(T, t)$, intervals in $\mathbb{R}$ have the standard metric, and $S^1 \times \Sigma_i^\pm$ and C have the metrics chosen above (which are independent of T, t), then the following composite maps should be isometries:

$$\begin{aligned} [0, \pm t_i] \times S^1 \times \Sigma_i^\pm &\xrightarrow{\xi_{\pm t_i} \times \text{Id}} [0, 1] \times S^1 \times \Sigma_i^\pm \xrightarrow{\rho_i^\pm} N_i^\pm & \text{if } \pm t_i \geq 1 \\ [0, T] \times C &\xrightarrow{\xi_T \times \text{Id}} [0, 1] \times C \xrightarrow{\rho_C} N_C. \end{aligned}$$

- $g(T, t)$ is independent of T outside N_C .
- $g(T, t)$ is independent of t_i outside $N_i^+ \cup N_i^-$.
- For $t_i \geq 1$, $g(T, t)$ is independent of t_i outside N_i^+ .
- For $t_i \leq -1$, $g(T, t)$ is independent of t_i outside N_i^- .
- $g(T, t)$ is on product form on $\mathbb{R}_+ \times Y$.

As $T \rightarrow \infty$ we obtain from $(W, g(T, t))$ the following Riemannian manifolds with tubular ends:

$$\begin{aligned} W_\infty^- &= W^- \cup_C (\mathbb{R}_- \times C) \\ V_{i,\infty} &= V_i \cup_{\partial V_i} (\mathbb{R}_+ \times \partial V_i) \end{aligned}$$

The metric on W_∞^- is independent of t , while the metric on $V_{i,\infty}$ depends only on t_i and is denoted $g_i(t_i)$.

6 Reducibles

Let $s_i \in H^2(W)$ be the Poincaré dual of $[S_i]$, and choose a $U(2)$ bundle $E \rightarrow W$ such that $c_1(E) = w + \sum_i s_i$ modulo torsion, where w is as in Theorem 2. For any integer k let $M_{T,t}(E, k)$ denote the moduli space of projectively $g(T, t)$ anti-self-dual connections A in E which are asymptotically trivial over the end $\mathbb{R}_+ \times Y$ and has “relative second Chern class” k , ie

$$\frac{1}{8\pi^2} \int_W \text{tr}(F_A^2) = k - \frac{1}{2} c_1(E)^2, \quad (6)$$

where F_A is the curvature of A . We wish to determine the reducible connections in $\bigcup_t M_{T,t}(E, k)$ when T is large and $k \leq 0$.

For any $v \in H^2(W)$ let $f_{T,v} : \mathbb{R}^{b^+} \rightarrow \mathbb{R}^{b^+}$ be the map defined in Subsection 4.2, using the family of metrics $g(T, t)$, $t \in \mathbb{R}^{b^+}$. If $v \cdot [\Sigma_i^\pm] \neq 0$ for all i

and both signs, then the zeros of $f_{T,v}$ form a bounded set. Restricting $f_{T,v}$ to a large sphere, of radius $r > 0$ say, we then obtain a map

$$f_{T,r,v} : S_r^{b^+-1} \rightarrow \mathbb{R}^{b^+} \setminus 0$$

and we define

$$\deg(v) = \deg(f_{T,r,v}/|f_{T,r,v}|). \quad (7)$$

By the homotopy invariance of the degree, the right hand side is independent of T and $r \gg 0$, so $\deg(v)$ is well-defined.

Lemma 1 *For $T > 0$ sufficiently large the following holds.*

- (i) *If $k < 0$ then there are no reducibles in $M_{T,t}(E, k)$ for any t .*
- (ii) *Suppose $M_{T,t}(E, 0)$ contains a reducible connection which respects a splitting $E = L_1 \oplus L_2$. Then modulo torsion the Chern class $c_1(L_1 \otimes \bar{L}_2)$ has the form $v = z - \sum_i \epsilon_i s_i$, where $z \in w + 2\mathcal{L}$, $z^2 = w^2$, $\epsilon_i = \pm 1$. Moreover, for such v we have*

$$\deg(v) = \epsilon_1 \epsilon_2 \dots \epsilon_{b^+}.$$

Proof. Fix $k \leq 0$. If $[A] \in M_{T,t}(E, k)$ respects the splitting $E = L_1 \oplus L_2$ and B is the induced connection in $K = L_1 \otimes \bar{L}_2$ then $\phi = \frac{1}{2\pi i} F_B$ is an anti-self-dual, closed L^2 form (with respect to $g(T, t)$) which represents $v = c_1(K)$. Recall from [10, Section 4] that

$$c_1(E)^2 = v^2 + 4k.$$

Now suppose $[A_n]$ is a reducible point in $M_{T(n),t(n)}(E, k)$ for $n = 1, 2, \dots$, where $T(n) \rightarrow \infty$ as $n \rightarrow \infty$. Since $\|\phi_n\|_{L^2}$ is independent of n , by the above equation, and $c_1(E) \cdot [\Sigma_i^\pm] = \mp 1$ for all i , $\{t(n)\}$ must be a bounded sequence. After passing to a subsequence we may therefore assume that $\{t(n)\}$ converges in $\mathbb{R}^{b^+}$, and that $\{\phi_n\}$ converges in C^∞ over compact subsets of both W_∞^- and $V_{i,\infty}$, with limits z' and z'_i , respectively.

Note that z', z'_i are harmonic L^2 forms and therefore decay exponentially over the ends (see [1]). We identify the cohomology class $[z']$ with the element $z \in H^2(W)$ which maps to $(0, [z'])$ under the isomorphism

$$H^2(W) \rightarrow H^2(W^+) \oplus \ker[H^2(W^-) \rightarrow H^2(C)], \quad (8)$$

where $W^+ = W \setminus \text{int } W^- \approx \cup_i V_i$. Then $z \in w + 2\mathcal{L}$, so $z^2 \leq w^2$ since w is extremal in $\mathcal{L}$. We also identify $[z'_i]$ with the class $z_i \in H^2(W)$ which maps

to $([z'_i], 0)$ in (8). Then $z_i = \alpha_i \sigma_i + \beta_i s_i$, where α_i is an even integer and β_i is an odd integer. Anti-self-duality implies $z_i^2 \leq 0$, so $z_i^2 = \alpha_i^2 - \beta_i^2 \leq -1$. We therefore get, for large n ,

$$w^2 - b^+ = c_1(E)^2 = v_n^2 + 4k \leq z^2 + \sum_i z_i^2 + 4k \leq w^2 - b^+ + 4k.$$

This gives $k \geq 0$, with equality only if $z^2 = w^2$ and $\alpha_i = 0$, $\beta_i = \pm 1$ for each i .

We will now compute the degree of $v = z - \sum_i \epsilon_i s_i$. Let $\mathcal{H}_{T,t}^+$ be the space of self-dual, closed, L^2 2-forms on W with respect to the metric $g(T, t)$, and let $\omega_{i,T,t}$, $i = 1, \dots, b^+$ be the basis for $\mathcal{H}_{T,t}^+$ constructed in Subsection 4.2, using the surfaces Σ_i . Note that as $T \rightarrow \infty$, $\omega_{i,T,t}$ converges in C^∞ to zero over compact subsets of W_∞^- (since the intersection form on $\ker[H^2(W^-) \rightarrow H^2(C)]$ is negative definite; see [1, Proposition 4.9]) and also over compact subsets of $V_{j,\infty}$ for $j \neq i$, and it converges over compact subsets of $V_{i,\infty}$ to some $g_i(t_i)$ self-dual form $\eta_i = \eta_i(t_i)$ uniquely determined by the properties

$$\int_{\Sigma_i} \eta_i \geq 1; \quad \int_{V_{i,\infty}} \eta_i \wedge \eta_i = 1.$$

Moreover, the convergence is uniform for $|t| \leq r$, so $f_{T,v}|_{D(r)}$ converges in C^0 to $f_{v,1} \times \dots \times f_{v,b^+}$ as $T \rightarrow \infty$, where

$$f_{v,i} : \mathbb{R} \rightarrow \mathbb{R}, \quad s \mapsto \int_{V_{i,\infty}} \eta_i(s) \wedge v.$$

Now,

$$\int_{\Sigma_i} \eta_i(s) \pm \int_{S_i} \eta_i(s) = \int_{\Sigma_i^\pm} \eta_i(s) \rightarrow 0 \quad \text{as } s \rightarrow \pm\infty.$$

Hence $\int_{S_i} \eta_i(s)$ is negative for $s \gg 0$ and positive for $s \ll 0$. It follows easily from this that $\deg(v) = \epsilon_1 \epsilon_2 \dots \epsilon_{b^+}$. $\square$

In the next section we will need to understand which instantons over W may restrict to reducible connections over open subsets. We record here the following basic result.

We say a connection A in an $\mathrm{SO}(3)$ bundle P is *reducible* if A is preserved by a non-trivial automorphism of P , and we say A is *s-reducible* if A is preserved by a non-trivial automorphism of P which lifts to $P \times_{\mathrm{Ad}} \mathrm{SU}(2)$, cfr. [10, Section 2]. We denote by $\mathfrak{g}_P$ the bundle of Lie algebras associated to P .

Lemma 2 ([7, 12]) *Let Z be a connected, oriented Riemannian 4-manifold and A an anti-self-dual connection in a principal $SO(3)$ bundle $P \rightarrow Z$. Suppose A is not flat, and that A is reducible over some non-empty open subset of Z . Then A respects some splitting $\mathfrak{g}_P = \lambda \oplus L$ where λ is a real line bundle and L a real 2-plane bundle. Moreover, A is s -reducible if and only if λ is trivial.*

Proof. Let S be the set of points $x \in Z$ such that A is reducible in some open neighbourhood of x . Then S is open, and non-empty by assumption. The proof of [7, Lemma 4.3.21] shows that S is also closed, hence $S = Z$. Therefore, A locally respects splittings of the form $\mathfrak{g}_P = \lambda \oplus L$, where F_A takes values in λ . Since $(d_A^* d_A + d_A d_A^*) F_A = 0$, unique continuation (see [11]) implies that F_A cannot vanish in any non-empty open subset of Z . It follows that A respects a global splitting. The last assertion of the lemma is left to the reader. $\square$

7 Cutting down parametrized moduli spaces

From now on we fix a large $T > 0$ such that the conclusions of Lemma 1 hold, and suppress T from notation.

For any projectively flat connection ρ in $E|_Y$ set

$$\mathcal{M}_\rho = \bigcup_{t \in \mathbb{R}^{b^+}} (M_t(E, \rho) \times \{t\}),$$

where $M_t(E, \rho)$ is the moduli space of projectively $g(t)$ anti-self-dual connections in E with limit ρ over the end $\mathbb{R}_+ \times Y$. We will now explain what we shall mean by “cutting down” $\mathcal{M}_\rho^*$ (or more generally, a subset of $\mathcal{M}_\rho^*$) according to a monomial $\kappa \prod_i x_i^{n_i}$, where $\kappa = z_1 \cdots z_J$, $z_j \in H_{d_j}(W^-; \mathbb{Z})$, $0 \leq d_j \leq 2$, and each n_i is a non-negative integer. Each x_i may be thought of as the point class in $H_0(W; \mathbb{Z})$, but the location of the point will depend on t_i . The cut down moduli space will be denoted

$$(\kappa \prod_i x_i^{n_i}) \cdot \mathcal{M}_\rho^*, \tag{9}$$

although it depends on various choices not reflected in the notation.

Roughly speaking, to cut down $\mathcal{M}_\rho^*$ according to κ we first restrict connections to W^- (where the metric does not vary), and then cut down by the product of the μ -classes of the z_i just as on a closed 4-manifold. As for the factor $x_i^{n_i}$, let $\mathcal{M}_{\pm t_i \geq r}^*$ be the part of $\mathcal{M}_\rho^*$ where $\pm t_i \geq r$, and let $r \gg 0$. To cut down $\mathcal{M}_{\pm t_i \geq r}^*$ according to $x_i^{n_i}$ we restrict connections to a tubular

neighbourhood of $\Sigma_i^\pm$ where the metric does not vary, and then cut down by the n_i 'th power of the μ -class of a point. In the intermediate region $|t_i| \leq r$ we interpolate by “moving base-points”, as we will explain in a moment.

We will now make this precise. To cut down by κ , choose disjoint, compact, codimension 0 submanifolds $U_j \subset W^-$ such that z_j is the image of a class $z'_j \in H_{d_j}(U_j)$. To ensure that irreducible instantons over W restrict to irreducible connections over U_j we also require that $H_1(U_j; \mathbb{Z}/2) \rightarrow H_1(W; \mathbb{Z}/2)$ be surjective. Let $\mathcal{B}(U_j)$ be the orbit space of L_1^p connections in $E|_{U_j}$ with a fixed central part, and $\mathcal{B}^* \subset \mathcal{B}$ the irreducible part. Choose a generic geometric representative $R_j \subset \mathcal{B}^*(U_j)$ for $\mu(z'_j) \in H^{4-d_j}(\mathcal{B}^*(U_j))$, (see [7, 12]). Set

$$Z_\kappa = \{([A], t) \in \mathcal{M}_\rho^* \mid [A|_{U_j}] \in R_j \text{ for } j = 1, \dots, J\}.$$

We now turn to the factor $x_i^{n_i}$. For each i choose disjoint closed subintervals $\{I_{i\nu}\}_{1 \leq \nu \leq n_i}$ of $[\frac{2}{3}, 1]$, each with non-empty interior, and set

$$B_{i\nu}^\pm = \rho_i^\pm(I_{i\nu} \times S^1 \times \Sigma_i^\pm),$$

where $\rho_i^\pm$ is as in Section 5. Choose disjoint compact, connected codimension 0 submanifolds $K_{i\nu} \subset W \setminus \bigcup_{j \neq i} U_j$ such that

- $B_{i\nu}^\pm \subset K_{i\nu}$,
- The sets $K_{i\nu}$ are mutually disjoint and also disjoint from the sets U_j ,
- $H_1(K_{i\nu}; \mathbb{Z}/2) \rightarrow H_1(W; \mathbb{Z}/2)$ is surjective.

When the perturbations of the ASD equations are sufficiently small then the restriction map

$$r_{i\nu} : \mathcal{M}_\rho^* \rightarrow \mathcal{B}^*(K_{i\nu})$$

is well-defined, as follows from Lemma 2 and a compactness argument. Also, when $r \gg 0$ there is a well-defined restriction map

$$r_{i\nu}^\pm : (\mathcal{M}_\rho^*)_{\pm t_i > r} \rightarrow \mathcal{B}^*(B_{i\nu}^\pm).$$

This follows by a compactness and unique continuation argument from the fact that there are no reducible projectively flat connections in $E|_{S^1 \times \Sigma_i^\pm}$ (since $c_1(E) \cdot [\Sigma_i^\pm] = \mp 1$).

Choose a base-point $b_{i\nu}^\pm \in B_{i\nu}^\pm$ and a smooth path $\gamma_{i\nu} : [0, 1] \rightarrow K_{i\nu}$ such that $\gamma_{i\nu}(\pm 1) = b_{i\nu}^\pm$. Let $\mathbb{E}_{i\nu} \rightarrow \mathcal{B}^*(K_{i\nu})$ and $\mathbb{E}_{i\nu}^\pm \rightarrow \mathcal{B}^*(B_{i\nu}^\pm)$ be the

complexifications of the natural 3-plane bundles associated to the base-points b_i^+ and $b_i^\pm$, respectively. By means of holonomy along $\gamma_{i\nu}$ we can identify $\mathbb{E}_{i\nu}$ with the corresponding bundle with base-point $b_{i\nu}^-$.

For each pair i, ν choose a generic pair of sections of $\mathbb{E}_{i\nu}$; pulling these back by $r_{i\nu}$ gives a pair $(s_{i\nu 1}^0, s_{i\nu 2}^0)$ of sections of the bundle $\mathbb{F}_{i\nu} = r_{i\nu}^*(\mathbb{E}_{i\nu})$ over $\mathcal{M}_\rho^*$. Choose also a generic pair of sections of $\mathbb{E}_{i\nu}^\pm$; this gives a pair $(s_{i\nu 1}^\pm, s_{i\nu 2}^\pm)$ of sections of $\mathbb{F}_{i\nu}^\pm = (r_{i\nu}^\pm)^*\mathbb{E}_{i\nu}^\pm$. Let $\beta : \mathbb{R} \rightarrow \mathbb{R}$ be a smooth function such that $\beta(s) = 0$ for $|s| \leq r + 1$ and $\beta(s) = 1$ for $|s| \geq r + 2$. Because of the canonical identification $\mathbb{F}_{i\nu} = \mathbb{F}_{i\nu}^\pm$ over $(\mathcal{M}_\rho^*)_{\pm t_i > r}$, it makes sense to define sections $(s_{i\nu 1}, s_{i\nu 2})$ of $\mathbb{F}_{i\nu}$ by

$$s_{i\nu j} = \beta(t_i)s_{i\nu j}^\pm + (1 - \beta(t_i))s_{i\nu j}^0 \quad \text{for } \pm t_i \geq 0.$$

Now let

$$Z_{i\nu} \subset \mathcal{M}_\rho^*$$

be the locus where $s_{i\nu 1}, s_{i\nu 2}$ are linearly dependent. As explained in [12], $Z_{i\nu}$ is a disjoint union of finitely many smooth submanifolds, such that the top stratum has codimension 4 and a natural orientation, and there is no stratum of codimension 5. We now define

$$(\kappa \prod_i x_i^{n_i}) \cdot \mathcal{M}_\rho^* = Z_\kappa \cap (\cap_{i\nu} Z_{i\nu}).$$

8 Main mechanism of proof

We will use the following characterization of the invariant h , see [10] for more details. For an oriented integral homology 3-sphere V let $\text{CF}^*(V)$ be the $\mathbb{Z}/8$ graded Floer cochain complex of V with rational coefficients. Let $\delta' \in \text{CF}^1(V)$ and the homomorphism $\delta : \text{CF}^4(V) \rightarrow \mathbb{Q}$ be defined by counting with sign the points in 0-dimensional moduli spaces $\check{M}(\alpha, \theta)$ and $\check{M}(\theta, \alpha)$ over $\mathbb{R} \times V$, respectively, where α is an irreducible flat $\text{SU}(2)$ connection and θ the trivial $\text{SU}(2)$ connection. Here $\check{M} = M/\mathbb{R}$. We denote by $\delta'_0 \in \text{HF}^1(V)$ and $\delta_0 : \text{HF}^4(V) \rightarrow \mathbb{Q}$ the corresponding data in cohomology. Then

- Either $\delta_0 = 0$ or $\delta'_0 = 0$
- $h(V) > 0$ if and only if $\delta_0 \neq 0$
- $h(V) < 0$ if and only if $\delta'_0 \neq 0$.

By Proposition 2 we may assume $h(Y) = 0$. Taking $V = \overline{Y}$ above we can therefore find a cochain $\alpha = \sum_j c_j \alpha_j \in \text{CF}^0(\overline{Y})$, where the α_j 's are

generators and the coefficients are rational, such that

$$\delta' + d\alpha = 0. \quad (10)$$

For any non-positive integer k set

$$\mathcal{M}_k = \bigcup_{t \in \mathbb{R}^{b^+}} (M_t(E, k) \times \{t\}).$$

Let $v \in H^2(W; \mathbb{Z})/\text{torsion}$ be any of the classes in Lemma 1 (ii). The zeros of the map

$$f_v : \mathbb{R}^{b^+} \rightarrow \mathbb{R}^{b^+}$$

defined in (4) are precisely the parameter values of t for which the L^2 $g(t)$ -harmonic form representing v is anti-self-dual. By making a small, t -independent perturbation to the family of metrics $g(t)$ in some ball in W where $g(t)$ is independent of t we can arrange that 0 is a regular value of f_v for each v . Then $\mathcal{M}_0$ will contain only finitely many reducible points. After making a local perturbation to the ASD equations near each reducible point we can then arrange that all reducibles points in $\mathcal{M}_0$ are regular (see Subsection 4.2).

Let $\mathcal{M}'_0 \subset \mathcal{M}_0$ be the result of removing from $\mathcal{M}_0$ a small neighbourhood about each reducible point, such that $\partial\mathcal{M}'_0$ is a disjoint union of complex projective spaces. For $k < 0$ set $\mathcal{M}'_k = \mathcal{M}_k$. (Recall that there are no reducibles in $\mathcal{M}_k$ when $k < 0$.) Let m be as in Theorem 2 and set $n = (|w^2| - m)/2 - 1$. Since

$$\dim M_t(E, k) = 8k - 2c_1(E)^2 - 3(b^+ + 1)$$

and $c_1(E)^2 = w^2 - b^+$, we have

$$\dim \mathcal{M}_k = 8k + 4n + 2m + 1.$$

Consider the formal linear combination of moduli spaces

$$\tilde{\mathcal{M}}_k = \mathcal{M}'_k + \sum_j c_j \mathcal{M}_{\rho_j},$$

where ρ_j is a projectively flat connection in $E|_Y$ representing α_j such that $\dim \mathcal{M}_{\rho_j} = \dim \mathcal{M}_k$.

Suppose the conclusion of Theorem 2 does not hold, ie that

$$\lceil g/2 \rceil + b^+ - 1 < (|w^2| - m)/4. \quad (11)$$

Set $q = \lceil g/2 \rceil + b^+ - 1$. Then (11) is equivalent to $2q \leq n$. Let $a \in H_2(X; \mathbb{Z})$ be as in the theorem and write

$$(x_1^2 - 4)^{\lceil g/2 \rceil} \prod_{i \geq 2} (x_i^2 - 4) = \sum_d P_d,$$

where P_d is a homogeneous polynomial in $x_1, \dots, x_{b^+}$ of degree $2d$. Cutting down as in the previous section we define

$$\widehat{\mathcal{M}} = \sum_{d=0}^q (a^m x^{n-2q} P_d) \cdot \tilde{\mathcal{M}}_{d-q},$$

which we regard as a formal linear combination of smooth, oriented 1-manifolds with boundary.

In the last two sections we will show that the number of boundary points of $\widehat{\mathcal{M}}$, counted with multiplicity, is non-zero, while the number of ends is zero. This contradiction will prove Theorem 2.

9 Boundary points of $\widehat{\mathcal{M}}$

Let $\mathcal{T} \subset H^2(W; \mathbb{Z})$ be the torsion subgroup (which we may identify with the torsion subgroup of $H^2(X; \mathbb{Z})$).

Proposition 5

$$\# \partial \widehat{\mathcal{M}} = \pm 2^{b^+ - 1 - 2n - m} |\mathcal{T}| \eta(\mathcal{L}, w, a, m),$$

which is non-zero by the hypotheses of Theorem 2.

Proof. The boundary points of $\widehat{\mathcal{M}}$ all lie in $\mathcal{M}'_0$, so

$$\# \partial \widehat{\mathcal{M}} = \#(a^m x^{n-2q} P_q \cdot \partial \mathcal{M}'_0) = \#(a^m x^n \cdot \partial \mathcal{M}'_0).$$

Now let Q be the component of $\partial \mathcal{M}'_0$ corresponding to some reducible point $D = ([A], t)$. Fix the ordering of the corresponding splitting $E = L_1 \oplus L_2$ and set $v = c_1(L_1) - c_1(L_2)$. If we cut down Q according to the monomial $a^m x^n$ and count points with signs using the complex orientation of Q then the result is

$$\#(a^m x^n \cdot Q) = \epsilon 2^{-2n-m} (v \cdot a)^m,$$

where $\epsilon = \pm 1$ depends only on m, n , see [10, Section 4]. If $c = c_1(E)$ then the complex orientation of Q compare with the boundary orientation inherited from $\mathcal{M}'_0$ by

$$\epsilon' \epsilon_{v,t} (-1)^{((v+c)/2)^2},$$

where $\epsilon' = \pm 1$ is independent of D and the ordering of L_1, L_2 , and $\epsilon_{v,t} = \pm 1$ is the sign of the determinant of $df_v(t)$, see Proposition 4. The last factor is taken from [4, Prop. 3.25] and accounts for whether the complex orientation of $\det(\delta_D)$ agrees with the orientation of the whole determinant line $\det(\delta)$.

As explained in [10, Section 4], there are precisely $|\mathcal{T}|$ reducible points in $\mathcal{M}_0$ for every pair $(\{v, -v\}, t)$ where $v \in H^2(W; \mathbb{R})$ is one of the classes in Lemma 1 and $f_v(t) = 0$. Summing up we find that

$$\begin{aligned} \#\partial(a^m x^n \cdot \mathcal{M}'_0) &= \pm \frac{1}{2} |\mathcal{T}| \sum_v \sum_{t \in f_v^{-1}(0)} \epsilon_{v,t} (-1)^{((v+c)/2)^2} 2^{-2n-m} (v \cdot a)^m \\ &= \pm |\mathcal{T}| 2^{-2n-m-1} \sum_v \deg(v) (-1)^{((v+c)/2)^2} (v \cdot a)^m. \end{aligned}$$

If $v = z - \sum_i \epsilon_i s_i$ as in Lemma 1 then $\deg(v) = \epsilon_1 \cdots \epsilon_{b+}$, and a simple computation shows that

$$(-1)^{((z+w)/2)^2} = (-1)^{((v+c)/2)^2} \epsilon_1 \cdots \epsilon_{b+}.$$

From this the Proposition follows immediately. $\square$

The reader may wish to check that $\epsilon_{v,t} (-1)^{((v+c)/2)^2} (v \cdot a)^m$ does indeed remain unchanged when v is replaced by $-v$.

10 Ends of $\widehat{\mathcal{M}}$

There are two kinds of ends in $\widehat{\mathcal{M}}$. The first kind arises from factorizations through flat connections over $\mathbb{R}_+ \times Y$. Because of our choice of “limiting cycle” over the end $\mathbb{R}_+ \times Y$ (ie condition (10)), the number of ends of this kind (counted with multiplicity) is zero. Indeed, by gluing theory (see [6]) there is a finite collection of projectively flat connections z_i in $E|_Y$ and for each i a finite subset $K_i \subset \mathcal{M}_{z_i}$ equipped with a function $K_i \rightarrow \mathbb{Z}$ indicating multiplicities, such that the ends of $\widehat{\mathcal{M}}$ can be identified with

$$\sum_i K_i \times \left(\check{M}(z_i, \theta) + \sum_j c_j \check{M}(z_i, \rho_j) \right).$$

Therefore,

$$\#\{\text{ends of } \widehat{\mathcal{M}}\} = \sum_i (\#K_i) \langle [z_i], \delta' + \partial\alpha \rangle = 0,$$

where $\alpha, \delta' \in CF_4(Y) = CF^1(\overline{Y})$.

The second type of ends in $\widehat{\mathcal{M}}$ arises from neck-stretching. More precisely, these ends correspond to sequences $\{(t(\nu), [A(\nu)])\}_{\nu=1,2,\dots}$ in $\widehat{\mathcal{M}}$ where $|t(\nu)| \rightarrow \infty$ as $\nu \rightarrow \infty$. The goal of the remainder of this section is to prove that the number of ends of this kind, counted with multiplicities, is zero. We will make use of a recent result by Muñoz [14] which we first explain.

Let Σ be a closed surface of genus g and $F \rightarrow \Sigma$ the non-trivial $\mathrm{SO}(3)$ bundle. Consider the affinely $\mathbb{Z}/8$ graded Floer cohomology group HF_g^* of the $\mathrm{SO}(3)$ bundle $\mathbf{F} = S^1 \times F \rightarrow S^1 \times \Sigma$. Choosing an extension $\mathbf{F}' \rightarrow D^2 \times \Sigma$ of $\mathbf{F}$ we can fix a $\mathbb{Z}/8$ -grading of HF_g^* by decreeing that the element $1 \in \mathrm{HF}_g^*$ defined by counting points in zero-dimensional moduli spaces in $\mathbf{F}'$ has degree 0. (This grading is compatible with the canonical $\mathbb{Z}/2$ grading defined in [10, Subsection 2.2].)

The pair-of-pants cobordism gives rise to a product

$$\mathrm{HF}_g^p \otimes \mathrm{HF}_g^q \rightarrow \mathrm{HF}_g^{p+q}$$

which makes HF_g^* a commutative ring with unit. Let τ be the degree 4 involution of HF_g^* defined by the class $[S^1] \in H^1(S^1 \times \Sigma; \mathbb{Z}/2)$. This involution respects the product in the sense that $x \cdot \tau(y) = \tau(x \cdot y)$ for $x, y \in \mathrm{HF}_g^*$. Therefore the ring structure descends to the $\mathbb{Z}/4$ -graded quotient $\widetilde{\mathrm{HF}}_g^* = \mathrm{HF}_g^* / \tau$.

Let

$$\Psi_g : \mathrm{Sym}(H_{\mathrm{even}}(\Sigma)) \otimes \Lambda(H_1(\Sigma)) \rightarrow \mathrm{HF}_g^*$$

be the invariant defined by the bundle $\mathbf{F}'$, as explained in [10]. Let $x \in H_0(\Sigma; \mathbb{Z})$ be the point class and $\gamma_1, \dots, \gamma_{2g} \in H_1(\Sigma; \mathbb{Z})$ a symplectic basis with $\gamma_j \cdot \gamma_{j+g} = 1$. Set $\alpha = 2\Psi_g([\Sigma])$, $\beta = -4\Psi_g(x)$, and $\gamma = -2 \sum_{j=1}^g \Psi_g(\gamma_j \gamma_{j+g})$, and let $\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}$ denote the images of these classes in $\widetilde{\mathrm{HF}}_g^*$.

The mapping class group of Σ acts on HF_g^* and $\widetilde{\mathrm{HF}}_g^*$ in a natural way. Muñoz shows that the invariant part of $\widetilde{\mathrm{HF}}_g^*$ is generated as a ring by $\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}$, and he gives a recursive description of the ideal of relations. The result we will use here is the following.

Proposition 6 (Muñoz)

- (i) $\prod_{j=1}^g (\tilde{\beta} + (-1)^j 8) \in \tilde{\gamma} \widetilde{\mathrm{HF}}_g^2$
- (ii) $(\beta^2 - 64)^{\lceil g/2 \rceil} \in \gamma \mathrm{HF}_g^2$.

Part (ii) follows from part (i), which is stated explicitly in the proof of [14, Proposition 20].

We will now show that the number of ends of $\widehat{\mathcal{M}}$ coming from neck-stretching is zero.

Let $(t_\nu, [A_\nu])$, $\nu = 1, 2, \dots$ be a sequence of points in $\mathcal{M}_r$ with $|t_\nu| \rightarrow \infty$ as $\nu \rightarrow \infty$. The transversality assumptions imply that, after passing to a subsequence, $t_{i,\nu}$ will stay bounded for all but one value of i .

Now fix i and $\pm t_i = \tau \gg 0$, so that W contains a large cylinder isometric to $[0, \pm t_i] \times S^1 \times \Sigma_i^\pm$, while the other t_j 's vary freely. Then the corresponding part $\widehat{\mathcal{M}}_{\pm t_i = \tau}$ of $\widehat{\mathcal{M}}$ is a finite number of points (with multiplicities) which by gluing theory (see [6]) can be identified with a product of instantons over $\mathbb{R}^2 \times \Sigma_i^\pm$ and over $W \setminus \Sigma_i^\pm$ (with tubular end metrics). In particular, if we set $E_i^\pm = E|_{S^1 \times \Sigma_i^\pm}$, which is the pull-back of the $U(2)$ bundle over $\Sigma_i^\pm$ with $c_1 = -1$, then

$$\#\widehat{\mathcal{M}}_{\pm t_i = \tau} = \phi_i^\pm \cdot \psi_i^\pm$$

for certain $\phi_i^\pm \in \text{CF}^*(E_i^\pm)$ (measuring instantons over $\mathbb{R}^2 \times \Sigma_i^\pm$) and $\psi_i^\pm \in \text{CF}_*(E_i^\pm)$.

If $i \geq 2$ then $\Sigma_i^\pm$ has genus 1. Thus $\text{CF}^*(E_i^\pm)$ has only two generators, in degrees differing by 4 (see Appendix B), and

$$\phi_i^\pm = \beta^2 - 64 = 0, \quad i \geq 2$$

on chain level. This is essential, because it implies not only that $\#\widehat{\mathcal{M}}_{\pm t_i = \tau} = 0$, but also that $\psi_1^\pm$ is closed. The point here is that to prove $d\psi_1^\pm = 0$ one must consider (suitably cut down) moduli spaces over $W \setminus \Sigma_1^\pm$ of dimension 1, and these moduli spaces may have ends where $t_j \rightarrow \pm\infty$ for some $j \geq 2$.

Now let $\Phi^\pm, \Psi^\pm$ be the Floer (co)homology classes of $\phi_1^\pm, \psi_1^\pm$, respectively. By Muñoz' result we can write

$$\Phi^\pm = (\beta^2 - 64)^{\lceil g/2 \rceil} = \gamma e$$

for some $e \in \text{HF}_g^*$, so

$$\#\widehat{\mathcal{M}}_{\pm t_1 = \tau} = \gamma e \cdot \Psi^\pm = e \cdot \gamma \Psi^\pm.$$

But $\gamma \Psi^\pm$ is a linear combination of Floer homology classes each of which can be defined by counting points in moduli spaces over $W \setminus \Sigma_1^\pm$ cut down according to a monomial of the kind $(\prod_{j=1}^J z_j)(\prod_{i=2}^{b^+} x_i^{n_i})$ where $z_j \in H_{d_j}(W^-; \mathbb{Z})$ and $d_1 = 1$. Since $H_1(W^-; \mathbb{Q}) = 0$ and we are using rational coefficients for the Floer homology groups, this implies that $\gamma \Psi^\pm = 0$, as we will explain in the next section.

Given this, we conclude that if $\tau \gg 0$ then $\#\widehat{\mathcal{M}}_{\pm t_i = \tau} = 0$ for all i , and Theorem 2 follows.

11 μ -classes of loops

In this section we will give a simple proof of the vanishing result used at the end of the previous section (ie $\gamma\Psi^\pm = 0$). This is essentially part of the general assertion that the Donaldson-Floer invariants of 4-manifolds with boundary discussed in [10, Subsection 2.3] are well-defined, in particular independent of the choice of geometric representatives for μ -classes (see [12]). One approach to this is to adapt the proof for closed 4-manifolds given in [5] and [12], but we prefer a more direct argument which also yields more precise information.

Because of the complexity of the proof of Theorem 2 we will consider the following simplified situation. Let X be an oriented Riemannian 4-manifold with one tubular end $\mathbb{R}_+ \times Y$, and let $E \rightarrow X$ be a principal $\mathrm{SO}(3)$ bundle such that $E|_Y$ is admissible in the sense of [2]. To make sure moduli spaces are orientable we assume E lifts to a $\mathrm{U}(2)$ bundle, ie that $w_2(E)$ has an integral lift.

If Y is a homology sphere we will assume there are no reducibles in the instanton moduli spaces in E considered below.

Set $X_0 = X \setminus (\mathbb{R}_+ \times Y)$ and let $\mathcal{B}$ be the orbit space of L_1^p connections in $E_0 = E|_{X_0}$ modulo *even* gauge transformations of class L_2^p . Here p should be an even integer greater than 4, and ‘even’ means that the gauge transformation should lift to $E \times_{\mathrm{Ad}} \mathrm{SU}(2)$. If $x_0 \in X_0$ is a base-point then over the irreducible part $\mathcal{B}^* \subset \mathcal{B}$ we have a natural principal $\mathrm{SO}(3)$ bundle $\mathbb{P} \rightarrow \mathcal{B}^*$, the base-point fibration at x_0 . To this principal bundle and the adjoint representations of $\mathrm{SO}(3)$ on itself, its Lie algebra, and its double cover $\mathrm{Spin}(3)$ we associate three fibre bundles over $\mathcal{B}^*$, which we denote $\mathbb{G}$, $\mathbb{E}$, and $\tilde{\mathbb{G}}$, respectively. Note that $\mathbb{G}$ is the bundle of fibrewise automorphisms of $\mathbb{P}$.

Now consider a smooth loop $\lambda : S^1 \rightarrow X_0$ based at x_0 . Holonomy along λ defines a smooth section h of $\mathbb{G}$. For any flat connection α in $E|_Y$ set

$$Z_\alpha = \{[A] \in M(E, \alpha) \mid h(A) = 1\}, \quad (12)$$

where $M(E, \alpha)$ is the instanton moduli space in E with flat limit α . After perturbing h by a homotopy we may assume each Z_α is transversely cut out, and we obtain a Floer cocycle

$$\phi = \phi(\lambda) = \sum_{\alpha} (\#Z_\alpha) \alpha,$$

where the sum is taken over all equivalence classes of irreducible flat connections α for which $\dim M(E, \alpha) = 3$. For rational coefficients the result we

wish to prove is that the cohomology class $[\phi]$ is a linear function of the homology class $[\lambda]$. However, we will actually prove a more precise statement, which involves $\text{Spin}(3)$ holonomy:

Up to isomorphism there are two spin structures on the pull-back bundle $\lambda^*E \rightarrow S^1$. Such a spin structure consists of a principal $\text{Spin}(3)$ bundle $\pi : Q \rightarrow S^1$ together with a bundle homomorphism $Q \rightarrow \lambda^*E$, see [13]. If A is any connection in E then there is a unique connection B in Q such that $\pi(B) = \lambda^*(A)$, and we can look at the holonomy of B . Because we are restricting to *even* gauge transformations, this gives a section $\tilde{h}$ of $\tilde{\mathbb{G}}$ which maps to h under the covering $\tilde{\mathbb{G}} \rightarrow \mathbb{G}$ and which depends on the choice of Q only up to an overall sign. Now, $\tilde{\mathbb{G}}$ is a bundle of Lie groups isomorphic to $\text{Spin}(3)$, which we can think of as the group of unit quaternions, and the imaginary part h_0 of $\tilde{h}$ is a section of the vector bundle $\mathbb{E}$. Replacing the condition $h(A) = 1$ in (12) by $\tilde{h}(A) = \pm 1$ and $h_0(A) = 0$, respectively, we get Floer cocycles $\psi^\pm, \psi_0$, and these satisfy the relations

$$\phi = \psi_+ + \psi_-, \quad \psi_0 = \psi_+ - \psi_-.$$

Let $\Phi, \Psi^\pm, \Psi_0 \in \text{HF}^*(Y; \mathbb{Z})$ be the cohomology classes of $\phi, \psi^\pm, \psi_0$, respectively. Note that Ψ_0 is independent of λ , and since $\mathbb{E}$ has odd rank we have $2\Psi_0 = 0$. Therefore,

$$\Psi_+ = \Psi_-, \quad \Phi = 2\Psi^\pm \quad \text{modulo 2-torsion.}$$

Proposition 7

- (i) *The subset $\{\Psi_+(\lambda), \Psi_-(\lambda)\} \subset \text{HF}^*(Y; \mathbb{Z})$ depends only on the class of λ in $H_1(X; \mathbb{Z})$.*
- (ii) *$\lambda \mapsto \Psi^\pm(\lambda)$ defines a homomorphism*

$$H_1(X; \mathbb{Z}) \rightarrow \text{HF}^*(Y; \mathbb{Z})/2\text{-torsion}.$$

The Proposition is easily deduced from the following two lemmas:

Lemma 3 *If $\lambda_1, \lambda_2 : S^1 \rightarrow X_0$ are loops based at x_0 then for any compatible spin structures on $\lambda_1^*E, \lambda_2^*E$, and $(\lambda_1 \circ \lambda_2)^*E$, one has*

$$\Psi_-(\lambda_1 \circ \lambda_2) = \Psi_-(\lambda_1) + \Psi_-(\lambda_2).$$

Proof of Lemma: Set $G = \text{Spin}(3)$ and let $S\mathbb{H}_0$ denote the unit sphere in the space $\mathbb{H}_0$ of pure quaternions. (We identify $\mathbb{H}_0$ with the Lie algebra of G .) Define a subset $V \subset G \times G$ by

$$V = \{(\exp(sx), \exp(tx)) \mid x \in S\mathbb{H}_0, s, t \in [0, \pi], s + t \geq \pi\}.$$

If we ignore the singular points $(1, -1), (-1, 1), (-1, -1)$, then V is a smooth, orientable 4-manifold with boundary, and

$$\partial V = (\{-1\} \times G) \cup (G \times \{-1\}) \cup D,$$

where $D = \{(g_1, g_2) \in G \times G \mid g_1 g_2 = -1\}$.

If M is any closed, oriented 3-manifold and $f_1, f_2 : M \rightarrow G$ smooth maps such that $(f_1, f_2) : M \rightarrow G \times G$ misses the three singular points of V and is transverse to both V and ∂V then

$$0 = \#\partial(f_1 \times f_2)^{-1}(V) = \epsilon_1 \deg(f_1) + \epsilon_2 \deg(f_2) - \epsilon_3 \deg(f_1 \cdot f_2)$$

for some constants $\epsilon_1, \epsilon_2, \epsilon_3 = \pm 1$. Taking $M = G$, $f_1 = \text{Id}$, $f_2 \equiv 1$ gives $\epsilon_3 = \epsilon_1$, and similarly $\epsilon_3 = \epsilon_2$.

Let $\tilde{\mathbb{G}} \otimes \tilde{\mathbb{G}}$ denote the fibrewise product of $\tilde{\mathbb{G}}$ with itself. Since V is Ad_G -invariant it defines a subset $W \subset \tilde{\mathbb{G}} \otimes \tilde{\mathbb{G}}$ which is a fibre bundle over $\mathcal{B}^*$ with fibre V . Now let $\tilde{h}_j$ be the section of $\tilde{\mathbb{G}}$ obtained from λ_j , and set

$$Z'_\alpha = \{[A] \in M(E, \alpha) \mid (\tilde{h}_1(A), \tilde{h}_2(A)) \in W\}.$$

Then the Floer cochain

$$b = \sum_{\dim M(E, \alpha)=2} (\#Z'_\alpha) \alpha$$

satisfies

$$db = \epsilon_1 \psi_-(\lambda_1) + \epsilon_2 \psi_-(\lambda_2) - \epsilon_3 \psi_-(\lambda_1 \circ \lambda_2)$$

with the same constants ϵ_j as above. $\square$

Lemma 4 *If $\lambda : S^1 \rightarrow X_0$ represents the zero class in $H_1(X; \mathbb{Z})$ then there is a spin structure on λ^*E for which $\Psi_-(\lambda) = 0$.*

Proof. There is a smooth, compact, oriented, connected 2-manifold Σ with one boundary component (which we identify with S^1), together with a smooth map $f : \Sigma \rightarrow X_0$ such that $f|_{\partial\Sigma} = \lambda$. Choose a spin structure on f^*E . It is easy to see that if $\mu : S^1 \rightarrow \Sigma$ is any loop based at $z \in \partial\Sigma$ then $\Psi_-(f \circ \mu)$ depends only on the class of μ in the fundamental group $\pi_1(\Sigma, z)$. Since the identity map $S^1 \rightarrow \partial\Sigma$ is a product of commutators in $\pi_1(\Sigma, z)$, it follows from Lemma 3 that $\Psi_-(\lambda) = 0$. $\square$

A Determinant line bundles

This appendix gives an account of the construction of determinant line bundles for families of Fredholm operators, taking care of some signs that many authors have overlooked.

Determinant line bundles for families of elliptic operators arise naturally in gauge theory and symplectic geometry in connection with orientations of moduli spaces, see [7, 9]. In general, if $\{T(x)\}_{x \in C}$ is a continuous family of Fredholm operators between two Banach spaces then the determinant line bundle $\det(T)$, as a set, is the disjoint union of all the determinants $\det(T(x))$ as x varies through C , see below. There is a natural collection of local trivializations of $\det(T)$ which one can attempt to use to make $\det(T)$ a topological line bundle over C . What many authors seem to have overlooked is that the overlap transformations are not in general continuous. We resolve this problem by dividing the set of local trivializations into two parts, thereby obtaining an “even” and an “odd” topology on $\det(T)$. The two topologies are interchanged by the involution of $\det(T)$ which is multiplication with $(-1)^{\dim(\ker(T(x)))}$ on the fibre $\det(T(x))$.

For the sake of simplicity, we will in the main text only consider the *odd* topology (this prevents a sign in Equation 5).

A.1 Determinant lines

We first study the determinant line of a single Fredholm operator.

For a finite dimensional vector space A we denote by $\det(A)$ the highest exterior power of A . If

$$0 \rightarrow A_0 \xrightarrow{\alpha_0} A_1 \xrightarrow{\alpha_1} A_2 \rightarrow 0$$

is an exact sequence of linear maps between finite dimensional vector spaces then there is a natural isomorphism

$$\det(A_0) \otimes \det(A_2) \xrightarrow{\cong} \det(A_1), \quad x_0 \otimes x_2 \mapsto \alpha_0(x_0) \wedge s(x_2), \quad (13)$$

where $s : A_2 \rightarrow A_1$ is any right inverse of α_1 . More generally, if

$$0 \rightarrow A_0 \xrightarrow{\alpha_0} A_1 \xrightarrow{\alpha_1} \dots \xrightarrow{\alpha_{r-1}} A_r \rightarrow 0$$

is an exact sequence of linear maps one gets a natural isomorphism

$$\bigotimes_{i \text{ even}} \det(A_i) \xrightarrow{\cong} \bigotimes_{i \text{ odd}} \det(A_i).$$

Now let V, W be Banach spaces over $\mathbb{F} = \mathbb{R}$ or $\mathbb{C}$, and $S : V \rightarrow W$ a Fredholm operator. The determinant line of S is by definition

$$\det(S) = \det(\ker(S)) \otimes (\det(\operatorname{coker}(S)))^*.$$

If $f : \mathbb{F}^n \rightarrow W$ is a linear map set

$$S_f : V \oplus \mathbb{F}^n \rightarrow W \oplus \mathbb{F}^n, \quad (v, z) \mapsto (Sv + fz, 0),$$

and let $S \oplus f : V \oplus \mathbb{F}^n \rightarrow W$ be the W -component of S_f . Then there is a natural exact sequence

$$0 \rightarrow \ker(S) \rightarrow \ker(S_f) \rightarrow \mathbb{F}^n \xrightarrow{\bar{f}} \operatorname{coker}(S) \rightarrow \operatorname{coker}(S_f) \rightarrow \mathbb{F}^n \rightarrow 0,$$

where $\bar{f}$ is the map f followed by the projection onto $\operatorname{coker}(S)$. This sequence gives rise to a natural isomorphism

$$\det(S) \xrightarrow{\sim} \det(S_f). \quad (14)$$

Next we consider a pair of linear maps $f_j : \mathbb{F}^{n_j} \rightarrow W$, $j = 1, 2$. Set

$$f_1 \oplus f_2 : \mathbb{F}^{n_1+n_2} = \mathbb{F}^{n_1} \oplus \mathbb{F}^{n_2} \rightarrow W, \quad (z_1, z_2) \mapsto f_1 z_1 + f_2 z_2.$$

Then we have a diagramme of isomorphisms

$$\begin{array}{ccc} \det(S) & \xrightarrow{\lambda_1} & \det(S_{f_1}) \\ \lambda_3 \downarrow & & \downarrow \lambda_2 \\ \det(S_{f_1 \oplus f_2}) & = & \det((S_{f_1})_{f_2}) \end{array} \quad (15)$$

where all maps except the bottom horizontal one are instances of (14). Consider the three maps

$$\mathbb{F}^{n_1} \rightarrow \operatorname{coker}(S), \quad \mathbb{F}^{n_2} \rightarrow \operatorname{coker}(S \oplus f_1), \quad \mathbb{F}^{n_1+n_2} \rightarrow \operatorname{coker}(S) \quad (16)$$

obtained from $f_1, f_2, f_1 \oplus f_2$ in the obvious way. Let K_1, K_2, K_{12} be the kernels of the maps in (16), and let L_1, L_2, L_{12} be the images of the same maps.

Lemma 5 $\lambda_3 = (-1)^{\dim(L_1) \dim(K_2)} \lambda_2 \lambda_1$.

We remark that the sign in the Lemma does not depend on the sign-convention in (13) (ie the order of $\alpha_0(x_0), s(x_2)$).

Proof. The proof is an exercise in understanding the definitions involved. We will merely indicate where the sign comes from. Note that there is a commutative diagramme

$$\begin{array}{ccccccc}
& & 0 & & 0 & & 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \rightarrow & K_1 & \rightarrow & \mathbb{F}^{n_1} & \rightarrow & L_1 \rightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \rightarrow & K_{12} & \rightarrow & \mathbb{F}^{n_1+n_2} & \rightarrow & L_{12} \rightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \rightarrow & K_2 & \rightarrow & \mathbb{F}^{n_2} & \rightarrow & L_2 \rightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
& & 0 & & 0 & & 0
\end{array} \tag{17}$$

where all rows and columns are exact. This defines two isomorphisms

$$\det(K_1) \otimes \det(L_1) \otimes \det(K_2) \otimes \det(L_2) \xrightarrow{\sim} \det(\mathbb{F}^{n_1+n_2}),$$

which differ by the factor $(-1)^{\dim(L_1)\dim(K_2)}$. One isomorphism uses the top and bottom horizontal exact sequences and then the middle vertical sequence. The other isomorphism uses the remaining three exact sequences in the diagramme.

The point is that when computing $\lambda_2\lambda_1$ one uses the first isomorphism, while λ_3 involves the other one. The remaining details are left to the reader. $\square$

A.2 Determinant line bundles

Let $\mathcal{B}(V, W)$ the Banach space of bounded operators from V to W , and $\text{Fred}(V, W) \subset \mathcal{B}(V, W)$ the open subset consisting of all Fredholm operators. If C is a space and $T : C \rightarrow \text{Fred}(V, W)$ a continuous map we define $\det(T)$ as a set by

$$\det(T) = \bigcup_{x \in C} \{x\} \times \det(T(x)).$$

Note that if $T(x)$ is surjective for every x then $\ker(T) \subset C \times V$, the union of all sets $\{x\} \times \ker(T(x))$, is a topological vector bundle over C , so in this case

$\det(T)$ has a natural topology. For a general T , we will topologize $\det(T)$ by essentially specifying a set of local trivializations. Since the surjective operators are open in $\mathcal{B}(V, W)$, every point in C has an open neighbourhood U for which there exists a linear map $f : \mathbb{F}^n \rightarrow W$ such that $T(x) \oplus f : V \oplus \mathbb{F}^n \rightarrow W$ is surjective for every $x \in U$. By definition,

$$\det(T_f(x)) = \det(T(x) \oplus f) \otimes \det(\mathbb{F}^n)^*,$$

hence $\det(T_f)|_U$ has a natural topology. Moreover, fibrewise application of (14) gives a bijection

$$\lambda_{U,f,n} : \det(T)|_U \rightarrow \det(T_f)|_U.$$

Proposition 8 (i) *If $\epsilon \in \mathbb{Z}/2$ then $\det(T)$ has a unique topology such that the projection $\det(T) \rightarrow C$ is continuous and all maps $\lambda_{U,f,n}$ with $n \equiv \epsilon \pmod{2}$ are homeomorphisms. The corresponding space $\det_\epsilon(T)$ is a topological line bundle over C .*

(ii) *The fibre preserving map*

$$\det_0(T) \rightarrow \det_1(T)$$

which is multiplication by $(-1)^{\dim(\ker(T(x)))}$ on the fibre $\det(T(x))$, is a homeomorphism.

In the main text we will, as already mentioned, take $\epsilon = 1$.

Proof. The only remaining issue is continuity of overlap transformations. Suppose $f_j : \mathbb{F}^{n_j} \rightarrow W$, $j = 1, 2$ are linear maps such that $T(x) \oplus f_j$ is surjective for all x . Then we have a diagramme of bijective maps

$$\begin{array}{ccccccc} \det(T_{f_1}) & \xleftarrow{\lambda_1} & \det(T) & = & \det(T) & \xrightarrow{\lambda_2} & \det(T_{f_2}) \\ \beta_1 \downarrow & & \downarrow & & \downarrow & & \downarrow \beta_2 \\ \det((T_{f_1})_{f_2}) & = & \det(T_{f_1 \oplus f_2}) & = & \det(T_{f_2 \oplus f_1}) & = & \det((T_{f_2})_{f_1}) \end{array}$$

where all maps marked with an arrow are instances of (14). The middle square obviously commutes. In the fibres over $x \in C$ the square to the left commutes up to multiplication with $(-1)^{n_2 \dim(\text{coker}(T(x)))}$, by Lemma 5, and similarly the square to the right commutes up to multiplication with $(-1)^{n_1 \dim(\text{coker}(T(x)))}$. It follows that

$$\lambda_2 \lambda_1^{-1} = (-1)^{(n_1 + n_2) \dim(\text{coker}(T(x)))} \beta_2^{-1} \beta_1.$$

Since $\beta_2^{-1}\beta_1$ is continuous, and

$$\text{index}(T(x)) = \dim(\ker(T(x))) - \dim(\text{coker}(T(x)))$$

is locally constant, the Proposition follows. $\square$

A.3 Isomorphisms between determinant line bundles

Suppose $g : C \rightarrow \mathcal{B}(\mathbb{F}^p, W)$ is continuous and $\epsilon_1, \epsilon_2 \in \mathbb{Z}/2$. We will define a canonical isomorphism of line bundles

$$\gamma : \det_{\epsilon_1}(T) \rightarrow \det_{\epsilon_2}(T_g).$$

If $n_j \equiv \epsilon_j \pmod{2}$, and $f_j : \mathbb{F}^{n_j} \rightarrow W$ are linear maps such that $T \oplus f_1$ and $T \oplus g \oplus f_2$ are both surjective in some open set $U \subset C$, then $\det(T_{f_1})$ and $\det(T_{g \oplus f_2})$ can both be identified with $\det(T_{g \oplus f_1 \oplus f_2})$ over U . This gives an isomorphism $\det(T)|_U \rightarrow \det(T_g)|_U$, which is easily seen to be independent of f_1, f_2 , hence patch together to give the desired isomorphism γ .

In explicit computations it may be useful to have γ expressed in terms of the bijective map

$$\gamma' : \det_{\epsilon_1}(T) \rightarrow \det_{\epsilon_2}(T_g)$$

defined in (14). Set

$$d_1 = \dim(\text{coker}(T)), \quad d_2 = \dim(\text{coker}(T \oplus g)),$$

regarded as functions on C . Then Lemma 5 and a simple computation gives:

Proposition 9 $\gamma = (-1)^{d_1\epsilon_1 + d_2\epsilon_2 + d_1p + d_1d_2 + d_2}\gamma'.$

Finally, we remark that the isomorphism γ is functorial in the sense that if $g_j : C \rightarrow \mathcal{B}(\mathbb{F}^{p_j}, W)$ is a continuous map for $j = 1, 2$ and if $\epsilon_1, \epsilon_2, \epsilon_3 \in \mathbb{Z}/2$ then the composition of the isomorphisms

$$\det_{\epsilon_1}(T) \rightarrow \det_{\epsilon_2}(T_{g_1}) \rightarrow \det_{\epsilon_3}(T_{g_1 \oplus g_2})$$

agrees with the isomorphism $\det_{\epsilon_1}(T) \rightarrow \det_{\epsilon_3}(T_{g_1 \oplus g_2})$.

B On projectively flat $U(2)$ connections over 3-manifolds

In this appendix we will prove a simple result which describes (modulo $SU(2)$ gauge equivalence) projectively flat $U(2)$ connections over a closed, oriented 3-manifold in terms of flat $SU(2)$ connections over the complement of a suitable link. The existence of such a correspondence was pointed out in [2, p198]. We will then apply this result to compute the Floer chain complex of a non-trivial $SO(3)$ bundle over the 3-torus.

Let Y be a closed, oriented 3-manifold, $E \rightarrow Y$ a rank 2 Hermitian vector bundle, and $L = \Lambda^2 E$ the determinant line bundle of E . Let s be a regular, smooth section of L . The zero-set of s is then a disjoint union $\gamma = \coprod_j \gamma_j$ of embedded circles. Let $N \approx D^2 \times \gamma$ be a closed tubular neighbourhood of γ , and let $V \subset Y$ denote the complement of $(\frac{1}{2}\text{int} D^2) \times \gamma$. By modifying s if necessary we may arrange that $|s| = 1$ in V . Let $\hat{A}$ be a unitary connection in L such that $\nabla_{\hat{A}} s = 0$ in V . Since $L|_N$ is trivial there is a unitary isomorphism $L|_N \approx L_0 \otimes L_0$ where $L_0 = N \times \mathbb{C}$. Let $\hat{A}_0$ be the connection in L_0 induced by $\hat{A}$. Of course, $F(\hat{A}) = 2F(\hat{A}_0)$ in N .

For each j choose a point $z_j \in \gamma_j$ and define $m_j = S^1 \times \{z_j\}$. Then

$$\text{Hol}_{m_j}(\hat{A}_0) = \exp \left(- \int_{D^2 \times z_j} F(\hat{A}_0) \right) = -1,$$

where the second equality is a basic fact from Chern-Weil theory.

Let M_1 denote the moduli space of projectively flat unitary connections in E which induces $\hat{A}$ in L , modulo automorphisms of E of determinant 1. If P is the $SO(3)$ bundle associated to E then M_1 can be identified with the moduli space $\mathcal{R}_S(P)$ of flat connections in P modulo *even* automorphisms of P , ie those that lift to $P \times_{\text{Ad}(SO(3))} SU(2)$. If $[A] \in M_1$ then A and $\hat{A}_0 \oplus \hat{A}_0$ restrict to gauge equivalent connections over $D^2 \times z_j$, hence $\text{Hol}_{m_j}(A) = -1$.

Let M_2 denote the moduli space of flat unitary connections in $E|_V$ satisfying $\nabla_A s = 0$ and $\text{Hol}_{m_j}(A) = -1$ for all j , modulo automorphisms of $E|_V$ of determinant 1.

Proposition 10 *The restriction map $r : M_1 \rightarrow M_2$ is a bijection.*

Proof. Given $[B] \in M_2$ it is easy to find a flat $SU(2)$ connection A_0 over N such that if $A = A_0 \otimes \hat{A}_0$ then A and B restrict to gauge equivalent $U(2)$ connections over $N \cap V$. Gluing these together along $N \cap V$ produces an element $\alpha \in M_1$ such that $r(\alpha) = [B]$. Hence r is surjective. It is easy to see that r is also injective. $\square$

Proposition 11 *If $P \rightarrow T^3$ is any non-trivial $SO(3)$ bundle then $\mathcal{R}_S(P)$ has exactly two elements. These are both non-degenerate and differ in index by 4.*

Proof. Recall that $SO(3)$ bundles over a compact 3-manifold (or over any finite CW-complex of dimension ≤ 3) are determined up to isomorphism by their second Stiefel-Whitney class. Choose an indivisible class $c \in H^2(T^3; \mathbb{Z})$ which is a lift of $w_2(P)$. Then there exists a diffeomorphism of T^3 (defined by some element of $SL(3, \mathbb{Z})$) such that f^*c is the Poincaré dual of $[S^1] \times 1 \times 1$. We may therefore assume that $P = S^1 \times P_0$, where $P_0 \rightarrow T^2$ is the non-trivial $SO(3)$ bundle. Applying Proposition 10 with $\gamma = S^1 \times \text{pt}$ we find that $\mathcal{R}_S(P)$ has exactly two elements $\alpha_{\pm}$, given by the representations of $\pi_1(S^1 \times (T^2 \setminus \text{pt}))$ into $SU(2) = \text{Sp}(1)$ which take the three standard generators to $\pm 1, i, j$, respectively, where $1, i, j, k$ is the usual basis for the quaternion algebra.

It is not hard to see that the corresponding flat connection in P is non-degenerate, by observing that it pulls back to the trivial connection under the 4-fold covering $T^3 \rightarrow T^3$, $(r, s, t) \mapsto (r, s^2, t^2)$. We know apriori that there is a degree 4 involution of $\mathcal{R}_S(P)$ (see [10]), so the index difference of $\alpha_{\pm}$ must be 4 mod 8. $\square$

References

- [1] M. F. Atiyah, V. K. Patodi, and I. M. Singer. Spectral asymmetry and Riemannian geometry. I. *Math. Proc. Camb. Phil. Soc.*, 77:43–69, 1975.
- [2] P. J. Braam and S. K. Donaldson. Floer’s work on instanton homology, knots and surgery. In H. Hofer, C. H. Taubes, A. Weinstein, and E. Zehnder, editors, *The Floer Memorial Volume*, pages 195–256. Birkhäuser, 1995.
- [3] S. K. Donaldson. An application of gauge theory to four dimensional topology. *J. Differential Geometry*, 18:279–315, 1983.
- [4] S. K. Donaldson. The orientation of Yang-Mills moduli spaces and 4-manifold topology. *J. Differential Geometry*, 26:397–428, 1987.
- [5] S. K. Donaldson. Polynomial invariants for smooth four-manifolds. *Topology*, 29:257–315, 1990.
- [6] S. K. Donaldson. *Floer Homology Groups in Yang-Mills Theory*. Cambridge University Press, 2002.

- [7] S. K. Donaldson and P. B. Kronheimer. *The Geometry of Four-Manifolds*. Oxford University Press, 1990.
- [8] A. Floer. An instanton invariant for 3-manifolds. *Comm. Math. Phys.*, 118:215–240, 1988.
- [9] A. Floer and H. Hofer. Coherent orientations for periodic orbit problems in symplectic geometry. *Math. Zeitschrift*, 212(1):13–38, 1993.
- [10] K. A. Frøyshov. Equivariant aspects of Yang-Mills Floer theory. *Topology*, 41:525–552, 2002.
- [11] J. Kazdan. Unique continuation in geometry. *Comm. Pure and Applied Math.*, 41:667–681, 1988.
- [12] P. B. Kronheimer and T. S. Mrowka. Embedded surfaces and the structure of Donaldson’s polynomial invariants. *J. Differential Geometry*, 41:573–734, 1995.
- [13] H. B. Lawson and M.-L. Michelsohn. *Spin Geometry*. Princeton University Press, 1989.
- [14] V. Muñoz. Ring structure of the Floer cohomology of $\Sigma \times S^1$. *Topology*, 38:517–528, 1999.

INSTITUT DES HAUTES ÉTUDES SCIENTIFIQUES,
F-91440 BURES-SUR-YVETTE, FRANCE